

1 Automatic Conjectures

These are the conjectures detected by the algorithm.

$$\frac{\pi}{-4} = \cfrac{1}{-1 + \cfrac{1}{-4 + \cfrac{-2}{-7 + \cfrac{-9}{-10 + \cfrac{-20}{-13 + \dots}}}}}$$

$$\frac{\pi}{-4} = \cfrac{1}{-1 + \cfrac{1}{-3 + \cfrac{4}{-5 + \cfrac{9}{-7 + \cfrac{16}{-9 + \dots}}}}}$$

$$\frac{1}{2} \left(\frac{\pi}{-2} + 1 \right) = \cfrac{1}{-3 + \cfrac{3}{-5 + \cfrac{8}{-7 + \cfrac{15}{-9 + \cfrac{24}{-11 + \dots}}}}}$$

$$\frac{\pi}{-2} + -1 = -3 + \cfrac{-2}{-6 + \cfrac{-9}{-9 + \cfrac{-20}{-12 + \cfrac{-35}{-15 + \dots}}}}$$

$$\frac{\pi}{-2} + 1 = \cfrac{1}{-2 + \cfrac{-1}{-5 + \cfrac{-6}{-8 + \cfrac{-15}{-11 + \cfrac{-28}{-14 + \dots}}}}}$$

$$\frac{\pi}{-2} = \cfrac{1}{-1 + \cfrac{-1}{-4 + \cfrac{-6}{-7 + \cfrac{-15}{-10 + \cfrac{-28}{-13 + \dots}}}}}$$

$$\frac{\pi}{-2} + -2 = \cfrac{1}{-1 + \cfrac{1}{2 + \cfrac{-2}{5 + \cfrac{-9}{8 + \cfrac{-20}{11 + \dots}}}}}$$

$$\frac{\pi}{-2} + -1 = \cfrac{1}{0 + \cfrac{1}{-3 + \cfrac{-2}{-6 + \cfrac{-9}{-9 + \cfrac{-20}{-12 + \dots}}}}}$$

$$-\left(\frac{\pi}{-2} + -1 \right) = 3 + \cfrac{-2}{6 + \cfrac{-9}{9 + \cfrac{-20}{12 + \cfrac{-35}{15 + \dots}}}}$$

$$\frac{1}{2}((-1) \cdot \pi + -2) = -3 + \frac{-2}{-6 + \frac{-9}{-9 + \frac{-20}{-12 + \frac{-35}{-15 + \dots}}}}$$

$$\frac{1}{-2}((-1) \cdot \pi + -2) = 3 + \frac{-2}{6 + \frac{-9}{9 + \frac{-20}{12 + \frac{-35}{15 + \dots}}}}$$

$$\frac{1}{-2}(\pi + 2) = -3 + \frac{-2}{-6 + \frac{-9}{-9 + \frac{-20}{-12 + \frac{-35}{-15 + \dots}}}}$$

$$\frac{1}{2}(\pi + 2) = 3 + \frac{-2}{6 + \frac{-9}{9 + \frac{-20}{12 + \frac{-35}{15 + \dots}}}}$$

$$-\left(\frac{\pi}{2} + 1\right) = -3 + \frac{-2}{-6 + \frac{-9}{-9 + \frac{-20}{-12 + \frac{-35}{-15 + \dots}}}}$$

$$\frac{\pi}{2} + 1 = 3 + \frac{-2}{6 + \frac{-9}{9 + \frac{-20}{12 + \frac{-35}{15 + \dots}}}}$$